

ISOLATED MPFL RECONSTRUCTION

# Physical Therapy, Strength and Conditioning

### PHASE I: MAXIMUM PROTECTION (WEEKS 0 TO 6)

· Weighting bearing as tolerated with brace locked in full extension for 6 weeks

### Week 0-2

- Range of motion 0-30 when not ambulating
- Elevate the ankle above the heart
- Multi-plane hip strengthening
- · Core and upper extremity strengthening

### Weeks 2-4

- Reduce inflammation
- Continue patellar mobilizations
- Range of motion 0-60 degrees when not ambulating
- Continue submaximal quadriceps activation
- Isometric hamstring/groin exercises

### Weeks 4-6

- Range of motion 0-90 degrees week 4-5, 0-120 degrees week 5-6 when not ambulating
- Begin straight leg raises
- Continue isometric hamstring/groin exercises
- Open chain hip strengthening

### PHASE II: PROGRESSIVE RANGE OF MOTION AND EARLY STRENGTHENING (WEEKS 6-12)

# Weeks 6-8:

- Open brace to 60° of flexion with weight bearing during week 6, 90° at week 7
- Continue with swelling control and patella mobility
- Gradually progress to full range of motion
- Normalize gait pattern
- Multi-plane straight leg raising and closed kinetic chain strengthening program
- Begin stationary bike
- Initiate shallow and deep water pool program

#### Weeks 8-10:

- Wean out of brace
- Normalize gait pattern
- Restore full ROM
- Gradual bilateral closed chain progression
- Gradual step up progression
- · Add light gym strengthening (leg press and hamstring curls)

### Weeks 10-12:

- Full ROM
- Terminal quadriceps stretching
- Advance closed chain and gym strengthening program
- Add treadmill walking and elliptical trainer

# PHASE III: PROGRESSIVE STRENGTHENING (WEEKS 12 TO 16)

- Full ROM
- Advance closed chain and gym strengthening
- Increase intensity on bike, treadmill, and elliptical trainer
- Initiate shallow water pool running program

# PHASE IV: ADVANCE STRENGTHENING, FUNCTIONAL DRILLS (WEEKS 16 TO 20)

- Begin linear land based running program
- Advance to lateral and rotational drills as tolerated

# PHASE V: PLYOMETRIC DRILLS, RETURN TO SPORT (WEEKS 20 TO 24)

- Advance gym strengthening
- Progress running/sprinting program
- Begin multi-directional field/court drills
- Begin bilateral progressing to unilateral plyometric drills
- Follow-up appointment with physician
- Sports test for return to competition **4.5 months post-op** per physician's release